

DKA diabetic ketoacidosis

Diabetic ketoacidosis (DKA)

Diabetic ketoacidosis (DKA) is a dangerous and potentially life threatening condition with thousands of preventable cases each year.


DKA most commonly happens in people with Type 1 diabetes, although it occasionally occurs in people with insulin-treated Type 2 diabetes. DKA happens when there is persistently high glucose in the blood and a lack of insulin.

There are two main stages of DKA.

Illness

Some types of illness can affect your diabetes control by raising your blood glucose levels. When you are ill, your body reacts by releasing more glucose into the bloodstream and increases insulin resistance, stopping it from working properly. This happens even if you are eating less food than usual or vomiting.

In someone without diabetes who is ill, the body simply releases more insulin to deal with the higher levels of glucose in the blood, bringing them back within the normal limits. However, in people with diabetes this is not done, increasing the risk of developing DKA.



Stage One

As the amount of glucose in the blood rises, the body tries to remove the excess by passing it out in the urine. This leads to dehydration, as the body moves water from the cells to the bloodstream to dilute the glucose. As this happens, levels of sodium and potassium (called electrolytes) are affected too. When these are unbalanced, you become even more poorly. Even though there is a lot of glucose in the blood, the lack of insulin means it can't get to the cells where it is needed for energy. The cells send out emergency signals and the body breaks down fat stores as replacement energy. As fat is broken down, poisonous acidic chemicals, called ketones, are released into the bloodstream.

Potential symptoms of stage one

- ✓ Blood glucose levels higher than 11.0 mmol/L
- ✓ Going to the toilet a lot to pass urine
- ✓ Thirst
- ✓ Tiredness
- ✓ Small amounts of ketones in the blood (0.6–1.5 mmol/L) or urine

If DKA is detected at stage one, with only small amounts of ketones in the body, it may be possible to bring things back to normal by taking extra short-acting insulin – your healthcare team will tell you how.

If you have Type 1 diabetes and you are pregnant, it is even more important that DKA does not progress, so seek medical advice straight away. You will probably be admitted to hospital to ensure the safety of both yourself and your baby.

Stage Two

As DKA progresses into its second stage, the amount of ketones in your blood rises. Ketones are poisonous and the body tries to remove them in the urine and on the breath. Higher levels of ketones in the blood can make you feel sick and can lead to vomiting. If untreated, high levels of ketones, dehydration and an imbalance of other chemicals in the blood can lead to unconsciousness and can eventually be fatal.

Potential symptoms of stage two

- ✓ Moderate to large amounts of ketones in the blood (over 1.5 mmol/L) or urine
- ✓ Breath smelling of pear drops
- ✓ Nausea and vomiting
- ✓ Drowsiness
- ✓ Deep rapid breathing
- ✓ Unconsciousness

If any of this happens, you will need immediate medical attention – usually being given fluids intravenously, and extra insulin to bring your blood glucose levels back under control. DKA can take up to 24 hours to develop in adults but develops faster in children.

It is important to get medical advice as soon as you think you may be developing DKA. Remember, at this stage DKA is a medical emergency and you should not try to treat it yourself.



Preventing DKA

Blood ketone testing

If you have Type 1 Diabetes or Type 2 Diabetes and being prescribed an SGLT-2i, it is important to test for ketones if your blood glucose levels are persistently over 11.1 mmol/L, or when you are ill. Some people do this by dipping a test strip into their urine, but others test for blood ketones using a meter.

What are the pros of testing for blood ketones?

- ✔ It gives you a here-and-now result, which means you can take action to bring your diabetes under control. The ketones in your urine were in your blood several hours ago
- ✔ It provides a numerical result rather than a colour comparison, which can be more accurate and easier to read. Blood testing, unlike urine testing, is unaffected by fluid intake
- ✔ If you find urine testing distasteful, blood testing is a quick and easy alternative

How can I test for blood ketones?

The test for blood ketones is the same process as for blood glucose.


If blood ketone testing is right for you then your GP can prescribe the appropriate strips.

Who is it most useful for?


- ✔ People with Type 1 diabetes
- ✔ Children, who are more likely to develop DKA quickly and who it might be difficult to collect urine from
- ✔ Insulin pumpers, as DKA can develop rapidly if the pump fails
- ✔ Pregnant women, who can also develop DKA quickly, which can seriously affect both their own health and that of their baby

Blood ketone testing doesn't replace blood glucose testing – it can be used alongside it. If you think it could help you manage your diabetes better, talk to your healthcare team.

Blood glucose test strip



Blood ketones test strip



What do the results mean?

<0.6

Under 0.6 mmol/L –

normal; consider rechecking blood ketone levels in 1-2 hours if blood glucose remains elevated, above 11 mmol/L.

0.6-1.5

0.6-1.5 mmol/L –

indicates a need for extra insulin. It is important the patient calls or follows the rules provided by their diabetes health care team and checks their blood glucose and blood ketone levels again in 1-2 hours.

>1.5

Over 1.5 mmol/L –

indicates risk of diabetic ketoacidosis. Patients should call their health care team **IMMEDIATELY**

When you're ill

Here are the basics to remember when you're poorly:

- ✓ Test blood glucose levels more often – at least four times a day – and test during the night as well
- ✓ If blood glucose levels are 11.1 mmol/L or more, test for blood ketones
- ✓ Continue to take insulin or diabetes medication and adjust the dose in response to test results if you have been taught to do so
- ✓ Drink plenty of sugar-free drinks
- ✓ If being sick, take carbohydrate-containing drinks such as milk and other milky drinks, fruit juice or sugary drinks such as Lucozade, ordinary cola or lemonade
- ✓ Eat little and often, taking carbohydrate containing drinks, as above, and snacks such as toast, biscuits and cereal

Contact your doctor or healthcare team if any of the following apply:

- ✓ Blood glucose levels are continuously high
- ✓ Ketones in blood or urine
- ✓ Vomiting
- ✓ Diarrhoea
- ✓ Or if you are unsure what to do




A.MENARINI
diagnostics

Diabetes Care Management Systems



**GlucoMen
areo GK**

**GLUCOFIX
TECH GK**



Glucoject®



Contactless download
to free diary and bolus
advice apps.

Blood glucose and ketone testing, lancets, pen needles and bolus advice

Why Menarini?

Menarini Diagnostics meters enable dual testing of blood glucose and blood ketone levels, for protection from DKA.

Always use your Menarini Diagnostics meter for both glucose and ketone testing to ensure that it is to hand and performing accurately and correctly at all times. It should not be used as a ketone only meter.

Symptoms of DKA usually evolve over a 24 hour period, with the first sign often being high blood glucose, so testing both glucose and ketones on the same meter will help ensure you are getting the best protection.

Call: (UK): 0800 243 667 (ROI): 1800 709 903

Email: customersupport@menarinidiag.co.uk

Visit: www.menarinidiag.co.uk

Follow:    